

# Where, in India, is hospital birth safer for babies than home birth?\*


scritto da Diane Coffey | 29 Aprile 2019



*L'Obiettivo di Sviluppo sostenibile 3 mira a ridurre la mortalità neonatale (NNM) a 12 decessi per 1000 nascite entro il 2030. Ridurre l'NNM in India è fondamentale per questo obiettivo perché questo paese conta il 27% dei decessi neonatali globali. Ricerche recenti, sintetizzate in questo articolo da Diane Coffey, suggeriscono che esiste una importante eterogeneità nell'associazione tra NNM e nascite ospedaliere nelle varie regioni dell'India.*

India contributes more neonatal deaths to global NNM (neonatal mortality) than any other country. Yet, India's national NNM of 30 per 1000 (IIPS and ICF 2017) masks wide spatial variations: among states with more than 25 million people, Uttar Pradesh had the highest NNM at 45 per 1000, and Kerala had the lowest at 4 per 1000. Figure 1 shows state level variation in NNM in India, and changes between 2005 and 2015. India's state-level variation in NNM is similar to the country-level variation that exists on a global scale: according to the 2015 World Development Indicators, the Central African Republic had the second highest NNM in the world at 43 per 1000, while the United States had an NNM of 4 per 1000 (World Bank Group 2015).

**Figure 1 - Neonatal mortality in India states, 2005 and 2015**



*Data sources: NFHS-3 and NFHS-4*

India's high NNM is surprising in light of a recent, dramatic increase in the fraction of births that occur in health facilities, rather than at home. In 2005, about 40% of births took place in health facilities; by 2015, this figure was 80%. This increase was in part due to a large-scale, conditional cash transfer program called *Janani Surksha Yojana* (JSY) launched in 2005 as part of the central government's new National Rural Health Mission (NRHM). JSY, which means Safe Motherhood Scheme, pays local health workers to accompany women to deliver in health facilities. Women who deliver in health facilities also receive a cash payment.

## **Health policy assumes that hospital birth is safer than home birth**

A key assumption of JSY, and of much of the Indian government's maternal and child health strategy, is that shifting births from homes to health facilities will reduce NNM (Rao 2017). However, this assumption has rarely been probed, and may not be correct in all times and places. For instance, in India's National Family Health Survey (NFHS)-3, collected before JSY was implemented in 2005, there was little difference in neonatal survival between babies born at home and those born in health facilities.


In new research using data on states with populations of more than 25 million from India's National Family Health Survey, 2015-2016 (NFHS-4), I seek to advance understanding of the association between neonatal survival and facility birth (Coffey, 2019). I find that, on average, in an uncontrolled

analysis, NNM was 12 per 1000 lower among children born in health facilities than among children born at home. However, there was important regional variation in the association between NNM and facility birth.

## The association between NNM and facility birth differs by region

Figure 2 plots associations between NNM and facility birth that control for a child's village (rural) or census block (urban), as well as a host of child, mother, and household-level demographic and socioeconomic characteristics (see Coffey, 2019). These controls are important because the uncontrolled association between NNM and facility birth may reflect omitted variables about the socioeconomic situation of the child's family or neighborhood, rather than a positive effect of being born in a hospital.

**Figure 2 - Associations between NNM and facility birth in India, 2015–2016**



Note: Figure 2, reprinted from Coffey (2019), plots OLS regression coefficients and confidence intervals from regressions of NNM on facility birth for regions and states of India. Gujarat and Maharashtra comprise “west;” Uttar Pradesh and Bihar comprise “focus;” Haryana, Punjab, and Rajasthan comprise “north;” Andhra Pradesh, Telangana, Karnataka, Kerala, and Tamil Nadu comprise “south;” Orissa, West Bengal, Jharkhand, and Assam comprise “east;” and Madhya Pradesh and Chhattisgarh comprise “central.” “not UP & Bihar” indicates a result that pools states in all other regions except Uttar Pradesh and Bihar.

Figure 2 shows that except in the “focus” states of Uttar Pradesh and Bihar, the association is negative, if not always statistically significant. Uttar Pradesh and Bihar are populous states bordering Nepal which have among the worst health and human development outcomes in the country. In east and central India, where outcomes were historically similar to those of Uttar Pradesh and Bihar, the association between neonatal death and facility birth is now both negative and statistically significant even after the introduction of this large set of controls.

Perhaps the most striking result presented in the Figure is found on the right side. In a pooled

regression using data from all of the regions except for Uttar Pradesh and Bihar, the association between neonatal death and facility birth is negative and statistically significant: NNM is about 7 per 1000 lower for births in a health facility than for home births. For Bihar, however, the relationship is slightly negative but not statistically significant. The point estimate for Uttar Pradesh is positive, and the confidence interval on the estimate does not overlap with that for the pooled estimate for the other regions.

## **Policy relevance and the need for further comparative research**

Considering that promoting facility birth has been the cornerstone of maternal and newborn health policy in India for much of the last decade, the finding that facility birth is uncorrelated with neonatal death in Uttar Pradesh and Bihar – which together contribute 43% of India's neonatal mortality – is concerning.

The lack of an association between neonatal death and facility birth coheres with prior qualitative research which suggests that the quality of maternal and newborn care in health facilities in this region is extremely poor (Jeffery and Jeffery 2010; Coffey 2014). It also coheres with the results of Semrau et al. (2017), who studied outcomes of a randomized controlled trial of the "Better Birth" coaching program, which aimed to improve the quality of maternal and newborn care in health facilities in 24 districts of Uttar Pradesh. The program consisted of 43 day-long coaching visits to each facility over a period of eight months. Nevertheless, twelve months after the program was implemented, researchers found only modest differences in the quality of maternal and newborn care provided in intervention vs. control facilities. Thus, both qualitative and quantitative research provide evidence that the government's rather singular focus on promoting facility birth may be misguided in this high-mortality region.

The robust, negative association between neonatal death and facility birth in the east and central regions is, however, encouraging, and suggests that the JSY program may have had heterogeneous effects. These regions saw much larger improvements in NNM in the last decade than were observed for Uttar Pradesh and Bihar. Robust associations between neonatal death and facility birth are consistent with evidence from the Million Deaths Study, which studied changes in causes of death between 2000 and 2015 in partnership with India's Sample Registration System (Fadel et al. 2017). The study found that declines in NNM came primarily from declines in birth trauma and birth asphyxia, which are causes of neonatal death that could plausibly be influenced by care at birth in a health facility. Future research should compare how delivery and postpartum care practices differ between the central and east regions and the states of Uttar Pradesh and Bihar.

\*"Articolo presente anche sul sito [www.niussp.org](http://www.niussp.org)"

## **References**

Coffey, Diane. (2019). The association between neonatal death and facility birth in regions of India. *Demographic Research*. 40: 417-430.

Coffey, D. (2014). Costs and consequences of a cash transfer for hospital births in a rural district of Uttar Pradesh, India. *Social Science & Medicine*, 114:89-96.

Fadel, S. A., Rasaily, R., Awasthi, S., Begum, R., Black, R. E., Gelband, H., Gerland, P., Kumar, R., Liu, L., Mathers, C., et al. (2017). Changes in cause-specific neonatal and 1- 59-month child mortality in India from 2000 to 2015: A nationally representative survey. *The Lancet*, 390(10106):1972-1980.

IIPS and ICF (2017). National family health survey (NFHS-4), 2015-16: India. Technical report, Mumbai: International Institute for Population Sciences.

Jeffery, P. and Jeffery, R. (2010). Only when the boat has started sinking: A maternal death in rural north India. *Social Science & Medicine*, 71(10):1711-1718.

Rao, S. (2017). *Do We Care? India's Health System*. New Delhi: Oxford University Press.

Semrau, K. E., Hirschhorn, L. R., Marx Delaney, M., Singh, V. P., Saurastri, R., Sharma, N., Tuller, D. E., Firestone, R., Lipsitz, S., Dhingra-Kumar, N., et al. (2017). Outcomes of a coaching-based WHO Safe Childbirth Checklist program in India. *New England Journal of Medicine*, 377(24):2313-2324.

World Bank Group (2015). *World Development Indicators 2015* [electronic resource]. Washington, D.C.: World Bank Publications.